

INFORMATION EXTRACTION OF HARVEST INDEX FOR WINTER WHEAT BASED ON EOS-MODIS DATA IN HUANGHUAHAI PLAIN IN CHINA

Jianqiang REN^{1,2}, Xingren LIU³, Zhongxin CHEN^{1,2,*}, Yunyan DU⁴, Huajun TANG^{1,2}

(1 Key Laboratory of Resources Remote-Sensing & Digital Agriculture, Ministry of Agriculture, Beijing 100081, China; 2 Institute of Agricultural Resources & Regional Planning, Chinese Academy of Agricultural Sciences, Beijing 100081, China; 3 Synthesis Research Center of Chinese Ecosystem Research Network & Key Laboratory of Ecosystem Network Observation and Modeling, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China; 4 The State Key Laboratory of Resources and Environmental Information System, Institute of Geographic Science and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China)

Jianqiang REN, Tel: +8610 8210-9615 ext. 172; E-mail: hebjqren1975@126.com

Yunyan DU (presenting author) Tel: +8610 6488-8973; E-mail: duyy@lreis.ac.cn

*Zhongxin CHEN (Corresponding author), Tel: +8610 8210-8684; E-mail: zxchen@mail.caas.net.cn

Address: Institute of Agricultural Resources & Regional Planning, Chinese Academy of Agricultural Sciences, No 12 Zhongguancun South Avenue, Beijing 100081, China

Abstract: Harvest Index (HI) of crop is one of the most important factors affecting the crop yield. Many researches had been done on it to increase the crop yield by agronomist and agronomy breeding expert. Generally speaking, the HI of one crop has certain stability in a small region for a period of time and has a better positive relationship with its yield. These relationships have been widely used in crop simulation and crop yield prediction and estimation using remote sensing data. Under many situations, the HI parameter is regarded as a constant through referring to the published literatures or getting from field observations. But in a larger region, there is a certain variance for different varieties of a same kind of crop. There is no doubt that regarding the HI as a constant in a larger region is not reasonable and can't meet the need of accuracy of crop simulation and yield prediction. So early getting the spatial information of crop harvest index is meaningful to improve the accuracy of crop simulation and crop yield prediction.

In this paper, the authors attempted to develop a new method to extract the winter wheat harvest index information depending on the 250m 10-day EOS/MODIS NDVI data. The process was below as follows:

(1) two new parameters were structured after analyzing the crop growth profile which is the curve of time series of NDVI and time and that the parameters can indirectly reflect the meaning of HI. The two parameter are HI_{NDVI_SUM} and HI_{NDVI_k} .

$$HI_{NDVI_SUM} = f\left(\frac{\sum NDVI_{post}}{\sum NDVI_{pre}}\right)$$
$$HI_{NDVI_k} = -f\left(\frac{\bar{k}_{post_descend}}{k_{pre_ascend}}\right)$$

Where $\sum NDVI_{post}$ is the accumulative total 10-day MVC NDVI after maximum and reflect dry matter accumulation of grain; $\sum NDVI_{pre}$ is the accumulative total 10-day MVC NDVI before maximum and reflect dry matter accumulation of stem and leaves; $\bar{k}_{post_descend}$ is mean descend speed of NDVI from the stage of anthesis to the stage of mature and reflect

the growth condition of grain filling; \bar{k}_{pre_ascend} is mean ascend speed of NDVI from the stage of reviving to the stage of anthesis and reflect the growth condition of stem and leaves from the reviving stage to the anthesis stage; the parameters such as $\sum NDVI_{post} / \sum NDVI_{pre}$ and $\bar{k}_{post_descend} / \bar{k}_{pre_ascend}$ indirectly reflect the meaning of definition of HI which is the ratio of seed mass to total above-ground plant mass. The negative sign was to change the value of $\bar{k}_{post_descend} / \bar{k}_{pre_ascend}$ from negative to a positive number by force.

(2) relationships were established between the two parameters and the relevant field measured HI at the same point.

(3) using the new parameter derived from the update 10-day NDVI data to predict the HI spatial information and validation was done comparing the predicted HI and the measured HI in the field.

(4) analyzing the accuracy of each parameter predicting the HI and confirming the better parameter for extracting the HI spatial information.

Finally, we selected the Hengshui City (including 11 counties) as our study region which is important main crop producing region in Huanghuaihai Plain in China. Field measured points were distributed symmetrically in each county in Hengshui City. In order to improve the quality of 10-day MVC MODIS-NDVI, especially to eliminate the cloud-contaminated data and abnormal data in the MODIS-NDVI series, the Savitzky-Golay filter was applied to smooth the 10-day NDVI data. The field measured HI data of the year 2004 and 2007 including 71 field survey points were used to establish the relationships between the new parameters and measured HI. The field measured HI of the year of 2008 including 46 field survey points were used to validate the accuracy of the predicted spatial HI information. At last, the mean relative error of predicted HI of the two parameters HI_{NDVI_k} and HI_{NDVI_SUM} is 3.60% and 2.14% respectively. The corresponding RMSE of predicted HI of the two parameters is 0.04 and 0.02 respectively. We think HI_{NDVI_SUM} was more effective to predict HI than the parameter HI_{NDVI_k} although the two parameters both can get good spatial HI information at regional level.

Key words: harvest index; winter wheat; MODIS; NDVI; Savitzky-Golay filter